

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 167 531 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
02.01.2002 Bulletin 2002/01

(51) Int Cl.7: **C12N 15/82, A01H 5/00,**
A01H 5/10

(21) Application number: **01202314.9**

(22) Date of filing: **15.06.2001**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **22.06.2000 US 213567 P**
13.10.2000 US 241215 P
13.10.2000 US 240014 P

(71) Applicant: **Monsanto Technology LLC**
St. Louis, Missouri 63167 (US)

(72) Inventors:
• **Behr, Carl F.**
Wildwood, MO 63005 (US)
• **Hironaka, Catherine**
Pleasanton, CA 94588 (US)
• **Heck, Gregory R.**
Crystal Lake Park, MO 63131 (US)
• **You, Jinsong**
Ballwin, MO 63021 (US)

(74) Representative: **Bosch, Henry**
Monsanto Services International SA Avenue de
Tervuren 270-272
1150 Bruxelles (BE)

(54) **Corn transformant PV-ZMGT32 (NK603) and compositions and methods for detection thereof**

(57) The present invention provides a DNA construct that confers tolerance to transgenic corn plant. Also provided are assays for detecting the presence of the

PV-ZMGT32(nk603) corn event based on the DNA sequence of the recombinant construct inserted into the corn genome and of genomic sequences flanking the insertion site.

EP 1 167 531 A1

Description**FIELD OF THE INVENTION**

[0001] The present invention relates to the field of plant molecular biology, specifically the invention relates to a DNA construct for conferring glyphosate tolerance to a plant. The invention more specifically relates to a glyphosate tolerant corn plant PV-ZMGT32(nk603) and to assays for detecting the presence of corn plant PV-ZMGT32(nk603) DNA in a sample and compositions thereof.

BACKGROUND OF THE INVENTION

[0002] This invention relates to the glyphosate herbicide tolerant corn (*Zea mays*) plant PV-ZMGT32(nk603) and to the DNA plant expression construct of corn plant PV-ZMGT32(nk603) and the detection of the transgene/genomic insertion region in corn PV-ZMGT32(nk603) and progeny thereof.

[0003] Corn is an important crop and is a primary food source in many areas of the world. The methods of biotechnology have been applied to corn for improvement of the agronomic traits and the quality of the product. One such agronomic trait is herbicide tolerance, in particular, tolerance to glyphosate herbicide. This trait in corn has been conferred by the expression of a transgene in the corn plants (U.S. Patent No. 6,040,497).

[0004] The expression of foreign genes in plants is known to be influenced by their chromosomal position, perhaps due to chromatin structure (*e.g.*, heterochromatin) or the proximity of transcriptional regulation elements (*e.g.*, enhancers) close to the integration site (Weising *et al.*, Ann. Rev. Genet 22:421-477, 1988). For this reason, it is often necessary to screen a large number of events in order to identify an event characterized by optimal expression of a introduced gene of interest. For example, it has been observed in plants and in other organisms that there may be a wide variation in levels of expression of introduced genes among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative expression of a transgene in various plant tissues, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct. For this reason, it is common to produce hundreds to thousands of different events and screen those events for a single event that has desired transgene expression levels and patterns for commercial purposes. An event that has desired levels or patterns of transgene expression is useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are well adapted to local growing conditions.

[0005] It would be advantageous to be able to detect the presence of a particular event in order to determine whether progeny of a sexual cross contain a transgene of interest. In addition, a method for detecting a particular event would be helpful for complying with regulations requiring the premarket approval and labeling of foods derived from recombinant crop plants, for example. It is possible to detect the presence of a transgene by any well known nucleic acid detection method such as the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct unless the DNA sequence of chromosomal DNA adjacent to the inserted DNA ("flanking DNA") is known. An event-specific PCR assay is discussed, for example, by Windels *et al.* (Med. Fac. Landbouww, Univ. Gent 64/5b:459-462, 1999), who identified glyphosate tolerant soybean event 40-3-2 by PCR using a primer set spanning the junction between the insert and flanking DNA, specifically one primer that included sequence from the insert and a second primer that included sequence from flanking DNA.

SUMMARY OF THE INVENTION

[0006] According to one aspect of the invention, a DNA construct is provided that when expressed in plant cells and plants confers tolerance to glyphosate herbicide. This invention relates preferably to the methods for producing and selecting a glyphosate tolerant monocot crop plant. The DNA construct consists of two transgene expression cassettes. The first expression cassette comprising a DNA molecule of a rice (*Oryza sativa*) actin 1 promoter and rice actin 1 intron operably joined to a DNA molecule encoding a chloroplast transit peptide sequence, operably connected to a DNA molecule encoding a glyphosate resistant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS), operably connected to a DNA molecule comprising a 3' transcriptional terminator. The second transgene expression cassette of the DNA construct comprising a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter, operably connected to a DNA molecule comprising a Hsp70 intron, operably connected to a DNA molecule encoding a chloroplast transit peptide sequence, operably connected to a DNA molecule encoding a glyphosate resistant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS), operably connected to a DNA molecule comprising a 3' transcriptional termi-

nator.

[0007] More specifically, a DNA construct is provided that when expressed in plant cells and plants confers tolerance to glyphosate herbicide. This invention relates preferably to the methods for producing and selecting a glyphosate tolerant corn plant. The DNA construct consists of two transgene expression cassettes. The first expression cassette consisting of a DNA molecule of a rice (*Oryza sativa*) actin 1 promoter and rice actin 1 intron operably joined to a DNA molecule encoding an *Arabidopsis* EPSPS chloroplast transit peptide sequence, operably connected to a DNA molecule encoding a glyphosate resistant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS) isolated from *Agrobacterium tumefaciens* sp. strain CP4, operably connected to a DNA molecule consisting of a nopaline synthase transcriptional terminator. The second transgene expression cassette consisting of a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter containing a tandem duplication of the enhancer region, operably connected to a DNA molecule consisting of a *Zea mays* Hsp70 intron, operably connected to a DNA molecule encoding an *Arabidopsis* EPSPS chloroplast transit peptide sequence, operably connected to a DNA molecule encoding a glyphosate resistant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS) isolated from *Agrobacterium tumefaciens* sp. strain CP4, operably connected to a DNA molecule consisting of a nopaline synthase transcriptional terminator.

[0008] According to another aspect of the invention, compositions and methods are provided for detecting the presence of the transgene/genomic insertion region from a novel corn plant designated PV-ZMGT32(nk603). DNA molecules are provided that comprise at least one junction sequence of PV-ZMGT32(nk603) selected from the group consisting of

5' TGTAGCGGCCACGCGTGG 3' (SEQ ID NO:9), 5' TACCACGCGACACACTTC 3' (SEQ ID NO:10), and 5' TGCT-GTTCTGCTGACTTT 3' (SEQ ID NO:11) and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site and is diagnostic for the event. The corn plant and seed comprising these molecules is an aspect of this invention.

[0009] A novel DNA molecule 5'ACCAAGCTTTTATAATAG 3' (SEQ ID NO:12) and the complement thereof, wherein this DNA molecule is novel in PV-ZMGT32(nk603) and its progeny, the corn plant and seed comprising this molecule are other aspects of this invention.

[0010] According to another aspect of the invention, DNA molecules that comprise the novel transgene/genomic insertion region, SEQ ID NO:7 and SEQ ID NO:8 and are homologous or complementary to SEQ ID NO:7 and SEQ ID NO:8 are an aspect of this invention.

[0011] DNA molecules that comprise a sufficient length of a transgene portion of the DNA sequence of SEQ ID NO:7 and a similar sufficient length of a 5' flanking corn DNA sequence of SEQ ID NO:7; or a similar sufficient length of a transgene portion of the DNA sequence of SEQ ID NO:8 and a similar sufficient length of a 3' DNA sequence flanking the transgene, wherein these DNA molecules are useful as DNA primers in DNA amplification methods so as to provide a DNA amplicon product specifically produced from PV-ZMGT32(nk603) DNA and its progeny are another aspect of the invention. DNA primers homologous or complementary to a length of SEQ ID NO:7 and SEQ ID NO:8 are an aspect of the invention. The amplicons produced using DNA primers that are diagnostic for corn event PV-ZMGT32(nk603) and its progeny are a subject of this invention.

[0012] According to another aspect of the invention, methods of detecting the presence of DNA corresponding to the corn event PV-ZMGT32(nk603) event in a sample are provided. Such methods comprise: (a) contacting the sample comprising DNA with a DNA primer set, that when used in a nucleic acid amplification reaction with genomic DNA extracted from corn event PV-ZMGT32 (nk603) produces an amplicon that is diagnostic for corn event PV-ZMGT32 (nk603); (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon. A pair of DNA molecules comprising a DNA primer set that are homologous or complementary to SEQ ID NO:7 or SEQ ID NO:8 that function in a nucleic acid amplification reaction to produce an amplicon DNA molecule diagnostic for PV-ZMGT32(nk603) are provided. More specifically, a pair of DNA molecules comprising a DNA primer set, wherein the DNA molecules are identified as SEQ ID NO:13 or complements thereof and SEQ ID NO:14 or complements thereof; SEQ ID NO:15 or complements thereof and SEQ ID NO:16 or complements thereof are provided. The amplicon comprising the DNA molecules of SEQ ID NO:13 and SEQ ID NO:14. The amplicon comprising the DNA molecules of SEQ ID NO:15 and SEQ ID NO:16. The amplicon produced by the afore described method can hybridize under stringent conditions to SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12.

[0013] According to another aspect of the invention, methods of detecting the presence of a DNA molecule corresponding to the PV-ZMGT32(nk603) event in a sample, such methods comprising: (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe molecule that hybridizes under stringent hybridization conditions with genomic DNA from corn event PV-ZMGT32(nk603) and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA. More specifically, a method for detecting the presence of a DNA molecule corresponding to the PV-ZMGT32(nk603) event in a sample, such methods, consisting of (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe molecule that consists of SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12, wherein said DNA probe molecule hybridizes under stringent hybridization

conditions with genomic DNA from corn event PV-ZMGT32(nk603) and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA.

[0014] According to another aspect of the invention, methods of producing a corn plant that tolerates application of glyphosate are provided that comprise the steps of: (a) sexually crossing a first parental corn line comprising the expression cassettes of the present invention, which confers tolerance to application of glyphosate, and a second parental corn line that lacks the glyphosate tolerance, thereby producing a plurality of progeny plants; and (b) selecting a progeny plant that tolerates application of glyphosate. Such methods may optionally comprise the further step of back-crossing the progeny plant to the second parental corn line to producing a true-breeding corn plant that tolerates application of glyphosate.

[0015] According to another aspect of the invention, a method is provided to select for glyphosate tolerant corn plants of the present invention and progeny thereof comprising extracting DNA from a plant sample, contacting a DNA with a marker nucleic acid molecule selected from the group consisting of SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12, or complements thereof, detecting the hybridization of said marker nucleic acid molecule to the DNA, and performing a marker assisted breeding analysis for the genetic linkage of the glyphosate tolerant trait to the marker nucleic acid molecule.

[0016] The present invention provides a method of producing a corn plant tolerant to glyphosate herbicide comprising transforming a corn cell with the DNA construct (pMON25496), selecting the corn cell for tolerance to the treatment with an effective dose of glyphosate, and growing the corn cell into a fertile corn plant. The fertile corn plant can be self pollinated or crossed with compatible corn varieties to produce glyphosate tolerant progeny.

[0017] The invention further relates to a DNA detection kit comprising at least one DNA molecule of sufficient length of contiguous nucleotides homologous or complementary to SEQ ID NO:7 or SEQ ID NO:8 that functions as a DNA primer or probe specific for corn event PV-ZMGT32(nk603) and/or its progeny.

[0018] This invention further relates to the plants and seeds of glyphosate tolerant corn (*Zea mays*) PV-ZMGT32 (nk603) having ATCC Accession No. PTA-2478 and the progeny derived thereof. The corn plant or its parts produced by growing of the glyphosate tolerant corn plant PV-ZMGT32(nk603), the pollen and ovules of the corn plant PV-ZMGT32(nk603). The nuclei of vegetative cells, the nuclei of pollen cells, and the nuclei of egg cells of the corn plant PV-ZMGT32 (nk603) and the progeny derived thereof. The corn plant and seed PV-ZMGT32(nk603) from which the DNA primer molecules of the present invention provide a specific amplicon product is an aspect of the invention.

[0019] The foregoing and other aspects of the invention will become more apparent from the following detailed description and accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] Figure 1. Plasmid map of pMON25496

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0021] The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger *et al.*, *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag: New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994. The nomenclature for DNA bases as set forth at 37 CFR § 1.822 is used.

[0022] As used herein, the term "corn" means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

[0023] As used herein, the term "comprising" means "including but not limited to".

[0024] "Glyphosate" refers to N-phosphonomethylglycine and its salts. Glyphosate is the active ingredient of Roundup® herbicide (Monsanto Co.). Treatments with "glyphosate herbicide" refer to treatments with the Roundup®, Roundup Ultra®, Roundup UltraMax® herbicide or any other herbicide formulation containing glyphosate. The selection of application rates for a glyphosate formulation that constitute a biologically effective dose is within the skill of the ordinary agricultural technician.

[0025] A DNA construct is an assembly of DNA molecules linked together that provide one or more expression cassettes. The DNA construct is preferably a plasmid that is enabled for self replication in a bacterial cell and contains various endonuclease enzyme restriction sites that are useful for introducing DNA molecules that provide functional genetic elements, *i.e.*, promoters, introns, leaders, coding sequences, 3' termination regions, among others. The expression cassettes contained within a DNA construct comprise the necessary genetic elements to provide transcription of a messenger RNA. The expression cassettes can be designed to express in prokaryote cells or eukaryotic cells.

The expression cassettes of the present invention are designed to express most preferably in plant cells. The combination of expression cassettes in transgenic plants can lead to unexpected phenotypes. In the present invention, the combination of the rice actin 1 promoter cassette and the CaMV 35S promoter cassette provided in a DNA construct (pMON25496) enhanced transgenic corn plant tolerance to glyphosate herbicide when compared to the single cassettes alone and to the commercial standard so-called GA21 that contains three copies of a rice actin 1 promoter cassette.

[0026] A transgenic "event" is produced by transformation of plant cells with heterologous DNA construct, including a nucleic acid expression cassette that comprises a transgene of interest, the regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. The term "event" refers to the original transformant and progeny of the transformant that include the heterologous DNA. The term "event" also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term "event" also refers to DNA from the original transformant comprising the inserted DNA and flanking genomic sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA. A glyphosate tolerant corn plant PV-ZMGT32 (nk603) can be bred by first sexually crossing a first parental corn plant consisting of a corn plant grown from the transgenic corn plant PV-ZMGT32 (nk603) having ATCC Accession No. PTA-2478 and progeny thereof derived from transformation with the expression cassettes of the present invention that tolerates application of glyphosate herbicide, and a second parental corn plant that lacks the tolerance to glyphosate herbicide, thereby producing a plurality of first progeny plants; and then selecting a first progeny plant that is tolerant to application of glyphosate herbicide; and selfing the first progeny plant, thereby producing a plurality of second progeny plants; and then selecting from the second progeny plants a glyphosate herbicide tolerant plant. These steps can further include the back-crossing of the first glyphosate tolerant progeny plant or the second glyphosate tolerant progeny plant to the second parental corn plant or a third parental corn plant, thereby producing a corn plant that tolerates the application of glyphosate herbicide.

[0027] It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison WI (1987).

[0028] A "probe" is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of genomic DNA from corn event PV-ZMGT32 (nk603) whether from a corn plant or from a sample that includes DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

[0029] "Primers" are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the present invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic-acid amplification methods.

[0030] Probes and primers are of sufficient nucleotide length to bind to the target DNA sequence specifically in the hybridization conditions or reaction conditions determined by the operator. This length may be of any length that is of sufficient length to be useful in the detection method of choice. Generally, 11 nucleotides or more in length, preferably 18 nucleotides or more, more preferably 24 nucleotides or more, and most preferably 30 nucleotides or more are used. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions. Preferably, probes and primers according to the present invention have complete DNA sequence similarity of contiguous nucleotides with the target sequence, although probes differing from the target DNA sequence and that retain the ability to hybridize to target DNA sequences may be designed by conventional methods.

[0031] Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook *et al.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989 (hereinafter, "Sambrook *et al.*, 1989"); *Current Protocols in Molecular Biology*, ed. Ausubel *et al.*, Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, "Ausubel *et al.*, 1992"); and Innis *et al.*, *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR-primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose

such as Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, MA).

[0032] Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, *e.g.*, by re-cloning and sequencing such sequences.

5 [0033] The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an
10 anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions. Similarly, the
15 molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook *et al.*, 1989, and by Haymes *et al.*, In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, DC (1985), Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order
20 for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

[0034] As used herein, a substantially homologous sequence is a nucleic acid molecule that will specifically hybridize to the complement of the nucleic acid molecule to which it is being compared under high stringency conditions. Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C, are known to those skilled in the art or can be found
25 in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50°C to a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied,
30 or either the temperature or the salt concentration may be held constant while the other variable is changed. In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 9, 10, 11 and 12 or complements thereof or fragments of either under moderately stringent conditions, for example at about 2.0 x SSC and about 65°C. In a particularly preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ
35 ID NO:9 through SEQ ID NO:12 or complements or fragments of either under high stringency conditions. In one aspect of the present invention, a preferred marker nucleic acid molecule of the present invention has the nucleic acid sequence set forth in SEQ ID NO:9 through SEQ ID NO:12 or complements thereof or fragments of either. In another aspect of the present invention, a preferred marker nucleic acid molecule of the present invention shares between 80% and 100% or 90% and 100% sequence identity with the nucleic acid sequence set forth in SEQ ID NO:9 through SEQ ID
40 NO:12 or complement thereof or fragments of either. In a further aspect of the present invention, a preferred marker nucleic acid molecule of the present invention shares between 95% and 100% sequence identity with the sequence set forth in SEQ ID NO:9 through SEQ ID NO:12 or complement thereof or fragments of either. SEQ ID NO:9 through SEQ IN NO:12 may be used as markers in plant breeding methods to identify the progeny of genetic crosses similar to the methods described for simple sequence repeat DNA marker analysis, in "DNA markers: Protocols, applications,
45 and overviews: (1997) 173-185, Cregan, et al., eds., Wiley-Liss NY; all of which is herein incorporated by reference in its' entirety. The hybridization of the probe to the target DNA molecule can be detected by any number of methods known to those skilled in the art, these can include, but are not limited to, fluorescent tags, radioactive tags, antibody based tags, and chemiluminescent tags.

[0035] Regarding the amplification of a target nucleic acid sequence (*e.g.*, by PCR) using a particular amplification primer pair, "stringent conditions" are conditions that permit the primer pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

[0036] The term "specific for (a target sequence)" indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

55 [0037] As used herein, "amplified DNA" or "amplicon" refers to the product of nucleic-acid amplification of a target nucleic acid sequence that is part of a nucleic acid template. For example, to determine whether the corn plant resulting from a sexual cross contains transgenic event genomic DNA from the corn plant of the present invention, DNA extracted from a corn plant tissue sample may be subjected to nucleic acid amplification method using a DNA primer pair that

includes a first primer derived from flanking sequence in the genome of the plant adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair, preferably plus about fifty nucleotide base pairs, more preferably plus about two hundred-fifty nucleotide base pairs, and even more preferably plus about four hundred-fifty nucleotide base pairs. Alternatively, a primer pair can be derived from flanking sequence on both sides of the inserted DNA so as to produce an amplicon that includes the entire insert nucleotide sequence (e.g., the MluI DNA fragment of the pMON25496 expression construct, Figure 1, approximately 6706 nucleotide base pairs). A member of a primer pair derived from the plant genomic sequence may be located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to the limits of the amplification reaction, or about twenty thousand nucleotide base pairs. The use of the term "amplicon" specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

[0038] Nucleic-acid amplification can be accomplished by any of the various nucleic-acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, *inter alia*, in U.S. Patent Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis *et al.*, Academic Press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 kb of genomic DNA and up to 42 kb of bacteriophage DNA (Cheng *et al.*, Proc. Natl. Acad. Sci. USA 91:5695-5699, 1994). These methods as well as other methods known in the art of DNA amplification may be used in the practice of the present invention. The sequence of the heterologous DNA insert or flanking DNA sequence from corn event PV-ZMGT32(nk603) can be verified (and corrected if necessary) by amplifying such sequences from DNA extracted from the ATCC deposit Accession No. PTA-2478 seed or plants using DNA primers derived from the sequences provided herein followed by standard DNA sequencing of the PCR amplicon or of the cloned DNA.

[0039] The amplicon produced by these methods may be detected by a plurality of techniques. One such method is Genetic Bit Analysis (Nikiforov, *et al.* Nucleic Acid Res. 22:4167-4175, 1994) where an DNA oligonucleotide is designed which overlaps both the adjacent flanking genomic DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking genomic sequence), a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labelled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0040] Another method is the pyrosequencing technique as described by Winge (Innov. Pharma. Tech. 00:18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent genomic DNA and insert DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking genomic sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

[0041] Fluorescence Polarization as described by Chen, *et al.*, (Genome Res. 9:492-498, 1999) is a method that can be used to detect the amplicon of the present invention. Using this method an oligonucleotide is designed which overlaps the genomic flanking and inserted DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking genomic DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0042] Taqman® (PE Applied Biosystems, Foster City, CA) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the genomic flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0043] Molecular Beacons have been described for use in sequence detection as described in Tyangi, *et al.* (Nature Biotech. 14:303-308, 1996) Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking genomic and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert

DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluorescent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0044] The following examples are included to demonstrate examples of certain preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLES

EXAMPLE 1

[0045] The PV-ZMGT32(nk603) (hence forth referred to as nk603) transgenic corn event was generated by micro-projectile bombardment of maize embryos (Songstad *et al.*, In Vitro Cell Plant 32:179-183, 1996) using a linear Mlu I DNA fragment derived from pMON25496 (Figure 1). This DNA fragment contains two transgene expression cassettes that collectively confer corn plant tolerance to glyphosate. The first cassette is composed of the rice actin 1 promoter and intron (P-Os.Act1 and I-Os.Act1, U.S. Patent No. 5,641,876), operably connected to an *Arabidopsis* EPSPS chloroplast transit peptide (TS-At.EPSPS:CTP2, Klee *et al.*, Mol. Gen. Genet. 210:47-442, 1987), operably connected to a glyphosate tolerant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS) from *Agrobacterium sp.* strain CP4 (AGRTU.aroA:CP4, U.S. Patent No. 5,633,435) and operably connected to a nopaline synthase transcriptional terminator (T-AGRTU.nos, Fraley *et al.* Proc. Natl. Acad. Sci. USA 80:4803-4807, 1983). The second transgene expression cassette consists of the cauliflower mosaic virus 35S promoter containing a tandem duplication of the enhancer region (P-CaMV.35S, Kay *et al.* Science 236:1299-1302, 1987; U.S. Patent No. 5,164,316), operably connected to a *Zea mays* Hsp70 intron (I-Zm.Hsp70, U.S. Patent No. 5,362,865), operably connected to an *Arabidopsis* EPSPS chloroplast transit peptide (TS-At.EPSPS:CTP2, Klee *et al.*, Mol. Gen. Genet. 210:47-442, 1987), operably connected to a glyphosate tolerant 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS) from *Agrobacterium sp.* strain CP4 (AGRTU.aroA:CP4, U.S. Patent No. 5,633,435) and operably connected to a nopaline synthase transcriptional terminator (T-AGRTU.nos, Fraley *et al.* Proc. Natl. Acad. Sci. USA 80:4803-4807, 1983). Post-bombardment, glyphosate-tolerant transgenic calli were selected on media containing 3 mM glyphosate and plants were subsequently regenerated. Three hundred-four plants from 91 independent transgenic events were produced, nk603 was selected from this population based on a superior combination of characteristics, including glyphosate tolerance, agronomic performance, and single transgenic insertion. Greenhouse and field evaluations of the nk603 event and its derived progeny indicated this transgenic insertion confers tolerance that exceeds commercial specifications for full vegetative and reproductive tolerance to 340 g glyphosate/acre (840 g glyphosate/hectare; 32 oz Roundup Ultra/acre) when applied at the V4 and V8 leaf stages.

EXAMPLE 2

[0046] The glyphosate tolerant nk603 corn event was compared to the current commercial standard, GA21 (U.S. Patent No. 6,040,497), for tolerance to glyphosate vegetative injury and effect on yield. GA21 contains at least 3 transgene expression cassettes arranged in tandem in the corn genome of GA21 event (SCP/GMO/232-Final, European Commission Health & Consumer Protection Directorate-General). The transgene cassette of GA21 consists of a rice actin 1 promoter and intron linked to a ribulose 1,5-bisphosphate carboxylase chloroplast transit peptide linked to a modified maize glyphosate resistant EPSPS and a 3' nopaline synthase transcription termination region. Plants of nk603 and GA21 were planted in rows in replicated field plots. The treatments were 1) unsprayed, 2) sprayed at a 64 ounces/acre rate of Roundup Ultra® at V4 leaf stage and again with 64 ounces/acre Roundup Ultra® at V8 leaf stage, 3) sprayed at a 96 ounces/acre rate of Roundup Ultra® at V4 leaf stage and again with 96 ounces/acre Roundup Ultra® at V8 leaf stage. Vegetative tolerance was measured as percent vegetative injury determined by the amount of leaf malformation observed 10 days after the herbicide treatment at the V8 leaf stage. The yield of each plot was measured in bushels/acre and the percent reduction in yield determined for each herbicide treatment relative to the unsprayed treatment. The results shown in Table 1 illustrates that nk603 shows a lower percent vegetative injury level than the GA21 plants and the observed percent yield reduction is also less for the nk603 event. A low incidence of vegetative injury was observed in the unsprayed plots, this observation is attributable to various environmental factors other than glyphosate herbicide exposure. The double expression cassette of pMON25496 in nk603 was compared to vegetative

injury and fertility rating of 3 independent corn events obtained from only the CaMV.35S promoter driving expression of the glyphosate tolerance gene (AGRTU.aroA:CP4). It was observed that the double expression cassette conferred a higher level of vegetative tolerance and reproductive tolerance than three independent corn events (ev 1, ev 2, and ev 3) containing only the expression cassette where the glyphosate tolerance gene expression was driven by the CaMV.35S promoter. A higher level of vegetative tolerance to glyphosate herbicide injury was observed for nk603 plus 3 additional corn events derived from pMON25496 when compared to the average injury of 6 corn events derived from a construct where the glyphosate tolerance gene expression was driven only by the rice actin promoter and intron (P-Os.Act1/I-Os.Act1). The plants transformed with the double expression cassette possessed a higher level of glyphosate tolerance to vegetative and fertility injury than plants derived from transformation with the single expression cassettes this resulted in improved resistance to yield loss due to glyphosate herbicide application. The pMON25496 construct provides two plant expression cassettes at a single location in nk603 that confers a higher level of glyphosate tolerance than the triple tandem insertion occurring in the commercial standard, GA21.

Table 1.

Glyphosate Tolerance of nk603 -Vegetative injury, Yield, and Fertility Rating				
Event	Treatment	% Veg injury*	Yield (bushels/acre)	% Yield red.
GA21	Unsprayed	0.3	142.2	
	64 oz Roundup Ultra® at V4 followed by 64 oz at V8	5.3	134.1	5.7
	96 oz Roundup Ultra® at V4 followed by 96 oz at V8	8.3	129.1	9.2
	Unsprayed	0.9	145.6	
	64 oz Roundup Ultra® at V4 followed by 64 oz at V8	2.9	138.5	4.9
	96 oz Roundup Ultra® at V4 followed by 96 oz at V8	4.7	140.1	3.8

	Treatment	Fertility Rating**
nk603	64 oz Roundup® Ultra at V8	4.5
CaMV.35S ev 1	64 oz Roundup® Ultra at V8	2.0
CaMV.35S ev 2	64 oz Roundup® Ultra at V8	2.2
CaMV.35S ev 3	64 oz Roundup® Ultra at V8	2.4

	Treatment	ave. % Veg injury*
nk603 plus 3 additional pMON25496 events	128 oz Roundup® Ultra at V4 followed by 128 oz at V8	22.9
Six P-Os.Act1 single cassette events	128 oz Roundup® Ultra at V4 followed by 128 oz at V8	28.9

* Veg injury is observed 10 days after V8 treatment is one measure taken to assess vegetative injury in response to glyphosate treatment. ** Male fertility rating: 4-5 = fully fertile; 3 = significantly reduced pollen shed; 0-2 = completely sterile - highly sterile, not suitable for commercial use.

EXAMPLE 3

[0047] The corresponding flanking DNA molecule from nk603 was cloned by using ligated adapters and nested PCR as described in the Genome Walkers kit (catalog # K1807-1, CloneTech Laboratories, Inc, Palo Alto, CA). First, genomic DNA from the nk603 event was purified by the CTAB purification method (Rogers *et al.*, Plant Mol. Biol. 5:69-76, 1985). The genomic DNA libraries for amplification were prepared according to manufacturer's instructions (Genome Walker™, CloneTech Laboratories, Inc, Palo Alto, CA). In separate reactions, genomic DNA was digested overnight at 37°C with the following blunt-end restriction endonucleases: *EcoRV*, *ScaI*, *DraI*, *PvuII*, and *StuI* (CloneTech Laboratories, Inc, Palo Alto, CA). The reaction mixtures were extracted with phenol:chloroform, the DNA was precipitated by the

addition of ethanol to the aqueous phase, pelleted by centrifugation, then resuspended in Tris-EDTA buffer (10mM Tris-.HCl, pH 8.0, 1mM EDTA). The purified blunt-ended genomic DNA fragments were ligated to the Genome Walker™ adapters according to the manufacturer's protocol. After ligation, each reaction was heat treated (70°C for 5 min) to terminate the reaction and then diluted 10-fold in Tris-EDTA buffer. One µl of each respective ligation was then amplified in a 50 µl reaction that included 1 µl of respective adapter-ligated library, 1 µl of 10 µM Genome Walker™ adapter primer AP1 (5'GTATATCGACTCACTATAGGGC 3', SEQ ID NO:1), 1 µl of 10 µM nk603 transgene-specific oligonucleotide

(5'TGACGTATCAAAGTACCGACAAAAACATCC 3' SEQ ID NO:2), 1 µl 10 mM deoxyribonucleotides, 2.5 µl dimethyl sulfoxide, 5 µl of 10X PCR buffer containing MgCl₂, 0.5 µl (2.5 units) of Amplitaq thermostable DNA polymerase (PE Applied Biosystems, Foster City, CA), and H₂O to 50 µl. The reactions were performed in a thermocycler using calculated temperature control and the following cycling conditions: 1 cycle of 95°C for 9 min; 7 cycles of 94°C for 2 s, 70°C for 3 min; 36 cycles of 94°C for 2 s, 65°C for 3 min; 1 cycle of 65°C for 4 min. One µl of each primary reaction was diluted 50-fold with water and amplified in a secondary reaction (1 µl of respective diluted primary reaction, 1 µl of 10 µM Genome Walker™ nested adapter primer AP2 (5'ACTATAGGGCAGCGTGGT 3', SEQ ID NO:3, supplied by manufacturer), 1 µl of 10 µM nk603 transgene-specific nested oligonucleotide

(5'CTTTGTTTTATTTTGGACTATCCCGACTC 3', SEQ ID NO:4), 1 µl 10 mM deoxyribonucleotides, 2.5 µl dimethyl sulfoxide, 5 µl of 10X PCR buffer containing MgCl₂, 0.5 µl (2.5 units) of Amplitaq thermostable DNA polymerase (PE Applied Biosystems, Foster City, CA), and H₂O to 50 µl] using the following cycling conditions: 1 cycle of 95°C for 9 min; 5 cycles of 94°C for 2 s, 70°C for 3 min; 24 cycles of 94°C for 2 s, 65°C for 3 min; 1 cycle of 65°C for 4 min.

[0048] PCR products, representing 5' regions that span the junction between the nk603 transgenic insertion and the neighboring flanking corn DNA were purified by agarose gel electrophoresis followed by purification from the agarose matrix using the QIAquick Gel Extraction Kit (catalog # 28704, Qiagen Inc., Valencia, CA) and direct cloning into the pGEM-T Easy vector (catalog # A1360, Promega, Madison, WI). The identity of the cloned PCR products and relationship to the Mlu I fragment of pMON25496 was confirmed by DNA sequence analysis (ABI Prism™ 377, PE Biosystems, Foster City, CA and DNASTAR sequence analysis software, DNASTAR Inc., Madison, WI).

[0049] Similarly, the nk603 3' flanking corn DNA sequence was amplified and cloned using nested gene specific primers, such as, SEQ ID NO:5

(5' AGATTGAATCCTGTTGCCGGTCTTGC 3') and SEQ ID NO:6

(5' GCGGTGTCATCTATGTTACTAGATCGGG 3') that anneal to the T-AGRTU.nos transcriptional terminator. Two T-AGRTU.nos transcriptional terminators are present in the nk603 transgenic/genomic insertion, one internal in the construct and one at the 3' end of the construct adjacent to corn genomic DNA. The PCR products produced in this reaction were sequenced and the DNA sequence that spans the junction between transgene and flanking sequence was distinguished from products of the internal T-AGRTU.nos by comparison to the known genetic element sequences of the pMON25496 construct as previously described.

[0050] Corn DNA sequence flanking both sides of the transgenic insertion was determined for nk603 by sequencing the Genome Walker™ -derived amplification products and alignment to known transgene sequence. At the 5' end of the transgenic insertion, the sequence of a 498 bp segment around the insertion junction was determined (SEQ ID NO:7). This consisted of 304 base pairs (bp) of the flanking maize genomic DNA sequence (nucleotides 1-304 of SEQ ID NO:7), 45 bp of pMON25496 construct DNA sequence (nucleotides 305-349 of SEQ ID NO:7) and 149 bp of DNA sequence from the 5' end of P-Os.Act1 (nucleotides 350-498 of SEQ ID NO:7).

[0051] The DNA sequence was determined for a 1183 bp segment around the 3' insertion junction (SEQ ID NO:8), that begins with 164 bp of the T-AGRTU.nos transcriptional terminator (nucleotides 1-164 of SEQ ID NO:8), 217 bp of pMON25496 construct DNA sequence (nucleotides 165-381 of SEQ ID NO:8), 305 bp of the maize plastid genes, *rps11* and *rpoA* (partial segments of each gene corresponding to bases 63-363 of Genbank accession X07810, corresponding to bases 382-686 of SEQ ID NO:8), and the remaining DNA sequence consisting of maize genomic DNA sequence flanking the integration site (corresponding to bases 687-1183 of SEQ ID NO:8).

[0052] The junction DNA molecules, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12 are novel DNA molecules in nk603 and are diagnostic for corn plant nk603 and its progeny. The junction molecules in SEQ ID NO:9, SEQ ID NO:10, and SEQ ID NO:11 represent about 9 nucleotides on each side of an insertion site of a transgene DNA fragment and corn genomic DNA. SEQ ID NO:9 is found at nucleotide positions 296-314 of SEQ ID NO:7. The junction sequences SEQ ID NO:10 and 11 are located at nucleotide positions 373-390 and 678-695, respectively, of SEQ ID NO:8, representing a junction DNA molecule of construct DNA sequence with corn plastid DNA sequence (SEQ ID NO:10) and construct sequence with corn genomic DNA sequence (SEQ ID NO:11). SEQ ID NO:12 is located at nucleotide position 156-173 of SEQ ID NO:8 and represents a novel DNA molecule in nk603 due to it being a fusion of the T-AGRTU.nos terminator sequence with an inverted fragment of the rice actin promoter DNA sequence.

EXAMPLE 4

[0053] DNA event primer pairs are used to produce an amplicon diagnostic for nk603. These event primer pairs include but are not limited to SEQ ID NO:13 and SEQ ID NO:14 for the 5' amplicon DNA molecule and SEQ ID NO:15 and SEQ ID NO:16 for the 3' amplicon DNA molecule. In addition to these primer pairs, any primer pair derived from SEQ ID NO:7 and SEQ ID NO:8 that when used in a DNA amplification reaction produces a DNA amplicon diagnostic for nk603 is an aspect of the present invention. The amplification conditions for this analysis is illustrated in Table 2 and Table 3 for the 5' transgene insert/genomic junction region. The same method is applied for amplification of the 3' amplicon DNA molecule using primer DNA molecules SEQ ID NO:15 and SEQ ID NO:16, however, any modification of these methods that use DNA molecules or complements thereof to produce an amplicon DNA molecule diagnostic for nk603 is within the ordinary skill of the art. In addition, a control primer pair (SEQ ID NO:17 and 18) for amplification of an endogenous corn gene is included as an internal standard for the reaction conditions. The analysis of nk603 plant tissue DNA extract sample should include a positive tissue DNA extract control from nk603, a negative DNA extract control from a corn plant that is not nk603, and a negative control that contains no template corn DNA extract. Additional DNA primer molecules of sufficient length can be selected from SEQ ID NO:7 and SEQ ID NO:8 by those skilled in the art of DNA amplification methods, and conditions optimized for the production of an amplicon that may differ from the methods shown in Table 2 and Table 3 but result in an amplicon diagnostic for nk603. The use of these DNA primer sequences with modifications to the methods of Table 2 and 3 are within the scope of the invention. The amplicon wherein at least one DNA primer molecule of sufficient length derived from SEQ ID NO:7 and SEQ ID NO:8 that is diagnostic for nk603 is an aspect of the invention. The amplicon wherein at least one DNA primer of sufficient length derived from any of the genetic elements of pMON25496 that is diagnostic for nk603 is an aspect of the invention. The assay for the nk603 amplicon can be performed by using a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler as shown in Table 3, or by methods and apparatus known to those skilled in the art

Table 2.

PCR procedure and reaction mixture for the confirmation of nk603 5' transgene insert/genomic junction region.			
Step	Reagent	Amount	Comments
1	Nuclease-free water	add to final volume of 20 μ l	-
2	10X reaction buffer (with $MgCl_2$)	2.0 μ l	1X final concentration of buffer, 1.5 mM final concentration of $MgCl_2$
3	10 mM solution of dATP, dCTP, dGTP, and dTTP	0.4 μ l	200 μ M final concentration of each dNTP
4	event primer (SEQ ID NO:13) (resuspended in 1X TE buffer or nuclease-free water to a concentration of 10 μ M)	0.4 μ l	0.2 μ M final concentration
5	event primer (SEQ ID NO:14) (resuspended in 1X TE buffer or nuclease-free water to a concentration of 10 μ M)	0.4 μ l	0.2 μ M final concentration
6	control primer (SEQ ID NO: 17) (resuspended in 1X TE buffer or nuclease-free water to a concentration of 10 μ M)	0.2 μ l	0.1 μ M final concentration
7	control primer (SEQ ID NO:18) (resuspended in 1X TE buffer or nuclease-free water to a concentration of 10 μ M)	0.2 μ l	0.1 μ M final concentration
8	RNase, DNase free (500 ng/ μ l)	0.1 μ l	50 ng/reaction
9	REDTaq DNA polymerase (1 unit/ μ l)	1.0 μ l (recommended to switch pipets prior to next step)	1 unit/reaction

Table 2. (continued)

PCR procedure and reaction mixture for the confirmation of nk603 5' transgene insert/genomic junction region.			
Step	Reagent	Amount	Comments
10	Extracted DNA (template): <ul style="list-style-type: none"> • Samples to be analyzed <ul style="list-style-type: none"> * individual leaves * pooled leaves (maximum of 50 leaves/pool) • Negative control • Negative control • Positive control 	<ul style="list-style-type: none"> • 10-200 ng of genomic DNA • 200 ng of genomic DNA • 50 ng of corn genomic DNA (notnk603) • no template DNA • 50 ng of nk603 genomic DNA 	-

Table 3. Suggested PCR parameters for different thermocyclers

Gently mix and, if needed (no hot top on thermocycler), add 1-2 drops of mineral oil on top of each reaction. Proceed with the PCR in a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler using the following cycling parameters.

Note: The MJ Engine or Eppendorf Mastercycler Gradient thermocycler should be run in the calculated mode. Run the Perkin-Elmer 9700 thermocycler with the ramp speed set at maximum.

Cycle No.	Settings: Stratagene Robocycler
1	94°C 3 minutes
38	94°C 1 minute 60°C 1 minute 72°C 1 minute and 30 seconds
1	72°C 10 minutes

Cycle No.	Settings: MJ Engine or Perkin-Elmer 9700
1	94°C 3 minutes
38	94°C 10 seconds 60°C 30 seconds 72°C 1 minute
1	72°C 10 minutes

Cycle No.	Settings: Eppendorf Mastercycler Gradient
1	94°C 3 minutes
38	94°C 15 seconds 60°C 15 seconds 72°C 1 minute
1	72°C 10 minutes

[0054] A deposit of the Monsanto Company, corn seed of event PV-ZMGT32(nk603) disclosed above has been made under the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110. The ATCC accession number is PTA-2478. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

[0055] Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

[0056] All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

5 <110> Monsanto Co
 Behr, Carl
 Hironaka, Catherine
 Heck, Gregory
 You, Jinsong

10 <120> Corn Event PV-ZMGT32(nk603) and Composition and Methods for Detection
 Thereof

 <130> 38-21(52258)B

15 <150> 60/213,567
 <151> 2000-06-22

 <150> 60/241,215
 <151> 2000-10-13

20 <150> 60/240,014
 <151> 2000-10-13

 <160> 16

25 <170> PatentIn version 3.0

 <210> 1
 <211> 22
 <212> DNA
 30 <213> artifical

 <220>
 <221> source
 <222> (1)..(22)
 35 <223> fully synthesized

 <400> 1
 gtatatcgac tcactatagg gc 22

40 <210> 2
 <211> 30
 <212> DNA
 <213> artifical

45 <220>
 <221> source
 <222> (1)..(30)
 <223> full synthesized

50 <400> 2
 tgacgtatca aagtaccgac aaaaacatcc 30

55 <210> 3

<211> 19
 <212> DNA
 <213> artificial
 5
 <220>
 <221> source
 <222> (1)..(19)
 <223> fully synthesized
 10
 <400> 3
 actatagggc acgcgtggt 19
 15
 <210> 4
 <211> 29
 <212> DNA
 <213> artificial
 20
 <220>
 <221> source
 <222> (1)..(29)
 <223> fully synthesized
 25
 <400> 4
 ctttggttta ttttggacta tcccgactc 29
 30
 <210> 5
 <211> 26
 <212> DNA
 <213> artificial
 35
 <220>
 <221> source
 <222> (1)..(26)
 <223> fully synthesized
 40
 <400> 5
 agattgaatc ctggtgcgg tcttgc 26
 45
 <210> 6
 <211> 28
 <212> DNA
 <213> artificial
 50
 <220>
 <221> source
 <222> (1)..(28)
 <223> fully synthesized
 55

<400> 6
 gcggtgtcat ctatgttact agatcggg 28

5

<210> 7
 <211> 498
 <212> DNA
 <213> artifical

10

<220>
 <221> source
 <222> (1)..(498)
 <223> 1-304 Zea maize genomic DNA
 305-349 construct vector DNA
 350-498 rice actin 1 promoter DNA

15

<400> 7
 aatcgatcca aaatcgcgac tgaaatgggtg gaagaaagag agaacagaga gcctcacgtt 60
 tccaggggtga agtatcagag gatttaccgc ccatgccttt tatggagaca agaaggggag 120
 gaggtaaaca gatcagcatc agcgcctcgaa agtttcgtca aaggatgcgg aactgtttcc 180
 25 agccgcgctc gccattcggc cagactcctc ctctctcggc atgagccgat cttttctctg 240
 gcattttccaa ccctagagac gtgcgtccct ggtgggctgc tcggccagca agccttgtag 300
 cggccccacgc gtggtaccaa gcttgatata cctagggcgg ccgcgttaac aagcttactc 360
 30 gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac aaaggtaaga 420
 ttaccgggtca aaagtgaaaa catcagttaa aagggtgtata aagtaaaata tcggtaataa 480
 35 aagggtggccc aaagtgaa 498

<210> 8
 <211> 1183
 <212> DNA
 <213> artifical

40

<220>
 <221> source
 <222> (1)..(1183)
 <223> 1-164 Agrobacterium tumefaciens nos 3' terminator
 165-381 construct vector DNA
 382-686 Zea maize plastid genes, rps11 and rpoA
 687-1183 Zea maize genomic DNA

45

<400> 8
 gacgttatatt atgagatggg tttttatgat tagagtcccg caattataca ttttaatacgc 60
 gatagaaaac aaaatatagc gcgcaaacta ggataaatta tcgcgcgcgg tgtcatctat 120

50

55

gttactagat cggggatata cccggggaat tcggtaccaa gcttttataa tagtagaaaa 180
 5 gagtaaattt cacttttggc caccttttat taccgatatt ttactttata ccacctttta 240
 actgatgttt tcacttttga ccaggtaatc ttacctttgt tttatttttg actatccga 300
 ctctcttctc aagcatatga atgacctcga gtaagcttgt taacgcggcc gccctagga 360
 10 tatcaagctt ggtaccacgc gacacacttc cactctagt tttgagtga tctgttata 420
 tcttctcgaa ccataacaga ctagtattat ttgatcattg aatcgtttat tctcttgaa 480
 15 agcggtttca ttttttttta cagacgtctt tttttaggag gtcgacatcc attatgcggc 540
 ataggtgtta catcgcgat acaacttaac cgtacaccac ttttagcaat ggctcgtaat 600
 gcggcatctc ttccgctacc agcacctttt accataactt ctgctcgttg caaaccact 660
 20 gtacgaatag catctactgc tgttctgctg actttatttt ttttaataaa gtgaaaaacc 720
 ataaaatgga caacaacacc ctgccttca ctaccggtcg gagcgacgcc gaagatgggg 780
 25 ttcaacacgg tcgcgacacg gatgcaacgg accctccaag ccaatactcg aggcgggacc 840
 gacgacgtag gcaggggtgg ccataacgac ggtggcggca tccaacttgt tctttccctt 900
 tctctgtctt caacttgccg cggcagtctg ctagaccag gggatgctgt gtggaggaga 960
 30 ggtcgcgggg cccgattttt atagcctggg cgaggacgag cttggccgaa ccgatccaga 1020
 gctctgcgca aatcacgaag aaccagtggg gccgctcgcg cctagccac cgccaggagc 1080
 ggggcttgtt gcgagccgta gcgtcgggaa ggggacgacc cgctaggggg gcccatgctc 1140
 35 cagcgcccag agagaaaaaa agaaaggaag gcgcgagatg atg 1183

<210> 9
 40 <211> 19
 <212> DNA
 <213> artifical

<220>
 45 <221> source
 <222> (1)..(19)
 <223> Zea maize genomic and vector DNA

50 <400> 9
 tgtagcggcc cacgcgtgg 19

<210> 10
 55 <211> 18

<212> DNA
 <213> artificial
 5 <220>
 <221> source
 <222> (1)..(18)
 <223> zea maize plastid DNA and vector DNA
 10 <400> 10
 taccacgcga cacacttc 18
 15 <210> 11
 <211> 18
 <212> DNA
 <213> artificial
 20 <220>
 <221> source
 <222> (1)..(18)
 <223> Zea maize genomic DNA and vector DNA
 25 <400> 11
 tgctgttctg ctgacttt 18
 30 <210> 12
 <211> 18
 <212> DNA
 <213> artificial
 35 <220>
 <221> source
 <222> (1)..(18)
 <223> Agrobacterium tumefaciens nos 3' terminator and rice actin promot
 er DN
 40 <400> 12
 accaagcttt tataatag 18
 45 <210> 13
 <211> 22
 <212> DNA
 <213> artificial
 50 <220>
 <221> source
 <222> (1)..(22)
 <223> fully synthesized
 55

<400> 13
aatcgatcca aaatcgcgac tg 22

5

<210> 14
<211> 22
<212> DNA
<213> artifical

10

<220>
<221> source
<222> (1)..(22)
<223> fully synthesized

15

<400> 14
ttcactttgg gccacctttt at 22

20

<210> 15
<211> 22
<212> DNA
<213> artifical

25

<220>
<221> source
<222> (1)..(22)
<223> fully synthesized

30

<400> 15
gacgttatatt atgagatggg tt 22

35

<210> 16
<211> 22
<212> DNA
<213> artifical

40

<220>
<221> source
<222> (1)..(22)
<223> fully synthesized

45

<400> 16
catcatctcg cgccttcctt tc 22

50 Claims

1. A DNA construct comprising: a first and a second expression cassette, wherein said first expression cassette in operable linkage comprises (i) a rice actin 1 promoter; (ii) a rice actin 1 intron; (iii) a chloroplast transit peptide encoding DNA molecule; (iv) a glyphosate tolerant EPSPS encoding DNA molecule; and (v) a transcriptional terminator DNA molecule; and said second expression cassette comprising in operable linkage (a) a CaMV 35S promoter; (b) a Hsp70 intron; (c) a chloroplast transit peptide encoding DNA molecule; (d) a glyphosate tolerant EPSPS encoding DNA molecule; and (e) a transcriptional terminator DNA molecule.

55

2. A DNA construct of claim 1, wherein the glyphosate tolerant EPSPS encoding DNA molecule consists of an AGRTU. *aroA*:CP4 DNA molecule.
3. A DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:7 or SEQ ID NO:8.
4. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of sufficient length of contiguous nucleotides of SEQ ID NO:7 or its complement to function as DNA primers or probes diagnostic for DNA extracted from corn plant PV-ZMGT32(nk603) or progeny thereof.
5. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of sufficient length of contiguous nucleotides of SEQ ID NO:8 or its complement to function as DNA primers or probes diagnostic for DNA extracted from corn plant PV-ZMGT32(nk603) or progeny thereof.
6. A method of detecting the presence of a DNA molecule selected from the group consisting of SEQ ID NO:7 and SEQ ID NO:8 in a DNA sample, the method comprising:
 - (a) extracting a DNA sample from a corn plant;
 - (b) contacting the DNA sample with a DNA primer pair comprising DNA primer molecules of sufficient length of contiguous nucleotides of SEQ ID NO:7 or its complement, or SEQ ID NO:8 or its complement;
 - (c) providing a nucleic acid amplification reaction condition;
 - (d) performing said nucleic acid amplification reaction, thereby producing a DNA amplicon molecule; and
 - (e) detecting the DNA amplicon molecule.
7. A method of detecting the presence of a DNA molecule selected from the group consisting of SEQ ID NO:7 and SEQ ID NO:8 in a DNA sample, the method comprising:
 - (a) extracting a DNA sample from a corn plant;
 - (b) contacting the DNA sample with a DNA molecule identified as SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12, wherein said DNA molecule is a DNA probe that hybridizes under stringent hybridization conditions with the DNA molecule selected from the group consisting of SEQ ID NO:7 or SEQ ID NO:8, and does not hybridize under the stringent hybridization conditions with a DNA sample not containing the DNA molecule identified as SEQ ID NO:7 or SEQ ID NO:8;
 - (c) subjecting the sample and probe to stringent hybridization conditions; and detecting hybridization of the probe to the DNA.
8. A DNA molecule selected from the group consisting of SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, and complements thereof.
9. A method of breeding a glyphosate tolerant trait into corn plants comprising
 - a) extracting a DNA sample from progeny corn plants;
 - b) contacting the DNA sample with a marker nucleic acid molecule selected from the group consisting of SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or complements thereof.
 - c) performing a marker assisted breeding method for the glyphosate tolerant trait, wherein the glyphosate tolerant trait is genetically linked to a complement of the marker nucleic acid molecule;
10. A method of producing a corn plant that tolerates application of glyphosate herbicide comprising:
 - (a) transforming a corn cell with the DNA construct of claim 1;
 - (b) selecting said corn cell for tolerance to application of glyphosate;
 - (c) growing said corn cell into a fertile corn plant.
11. A DNA detection kit comprising: at least one DNA molecule of sufficient length of contiguous nucleotides homologous or complementary to SEQ ID NO:7 or SEQ ID NO:8 that functions as a DNA primer or probe specific for corn event PV-ZMGT32(nk603) and/or its progeny.
12. A glyphosate tolerant corn plant, its progeny, or its parts thereof, said corn plant provided by ATCC seed deposit PTA-2478.




FIG. 1



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 01 20 2314 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	US 5 948 956 A (BERG JOHN BRADLEY ET AL) 7 September 1999 (1999-09-07) * column 10, line 11 - line 26; figure 4 * * column 10, line 55 - column 12, line 5 * ---	1,2,10	C12N15/82 A01H5/00 A01H5/10
Y,D	WINDELS P ET AL: "Development of a line specific GMO detection method: A case study." MEDEDELINGEN FACULTEIT LANDBOUWKUNDIGE EN TOEGEPASTE BIOLOGISCHE, vol. 64, no. 5b, 1999, pages 459-462, XP001032975 Thirteenth Forum for Applied Biotechnology.;Gent, Belgium; September 22-23, 1999, 1999 * the whole document * ---	3-8,11	
Y,D	WO 98 44140 A (DEKALB GENETICS CORP) 8 October 1998 (1998-10-08) * page 50, line 21 - line 26 * * page 63, line 20 - line 23 * * page 41, line 17 - page 46, line 20 * -----	3-8,11	
A		1,2,10	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N
INCOMPLETE SEARCH The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims. Claims searched completely : Claims searched incompletely : Claims not searched : 12 Reason for the limitation of the search: Article 53 (b) EPC - Plant variety			
Place of search		Date of completion of the search	Examiner
BERLIN		16 October 2001	De Kok, A
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 03-82 (P04C07)



European Patent
Office

Application Number

EP 01 20 2314

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☒ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

**LACK OF UNITY OF INVENTION
SHEET B**

Application Number
EP 01 20 2314

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1, 2, 10

A DNA construct comprising a first and second expression cassette, wherein the first expression cassette comprises a glyphosate tolerant EPSPS under the control of a rice actin-1 promoter and the second expression cassette comprises a glyphosate tolerant EPSPS under the control of a CaMV 34S promoter

2. Claims: 3-9 11 12

A glyphosate tolerant corn plant, PV-ZMGT32(nk206), ATCC seed deposit PTA-2478 and DNA molecules for the specific detection of said plant.

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 20 2314

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-10-2001

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 5948956	A	07-09-1999	AU	9806698 A	03-05-1999
			WO	9919499 A1	22-04-1999

WO 9844140	A	08-10-1998	US	6040497 A	21-03-2000
			AU	6882298 A	22-10-1998
			BR	9808475 A	23-05-2000
			EP	0975778 A1	02-02-2000
			WO	9844140 A1	08-10-1998

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82